

# There's an easier way to remove nutgrass and mullumbimby couch in turf.

## Sempra usage card

### Weeds controlled

- Nutgrass
- Mullumbimby Couch

### Turf Species

Sempra is safe to apply on a range of turf species if applied as per label directions

### When To Spray

**Nutgrass:** apply when new leaf growth is greater than 5cm

**Mullumbimby Couch:** apply when new leaf growth is greater than 2cm

- Apply when weeds are actively growing
- Apply follow-up treatments as required if sufficient new growth warrants treatment

### Weed Control Symptoms

- Weed foliage gradually turn yellow until total desiccation occurs.
- Initial symptoms appear in 7 - 10 days with full weed control effects appearing over 4 to 6 weeks.

### Sempra Application Rate

**Boomspray:** Sempra 65 - 130 g / ha plus 200mL / 100L water of a 600g / L non-ionic surfactant

**Spot Spraying:** Sempra 1.3 g/ 100 square metres plus 20mL / 10L water of a 600g / L non-ionic surfactant

**N.B. An adjuvant must be used when applying Sempra**

### How To Spray - Boomspray

**Rate:** 65 - 130 g / ha

**Water Volume:** apply in at least 80L / ha minimum water volume

**Nozzle Type:** flat fan nozzles

### How To Spray - Spot Spraying

**Equipment:** knapsack spray or handgun

Rate 1.3 g/ 100 square metres

**Water Volume:** mix rate in a volume of water eg. 10 litres that will treat 100 square metres



## Mixing

- Part fill spray tank
- Add Sempra gradually while under agitation
- Add the surfactant near the end of the spray fill process to avoid excessive foaming
- Use mixed spray solution within one day



## Critical Comments

To maximise the performance of spray application.

- Apply higher rates for dense weed conditions
- If an adjuvant is not used poor weed control will occur
- Do not apply after frosts have occurred
- Irrigation or rainfall within 2 hours of application will reduce control
- Do not mow for at least 2 days after treatment to ensure optimum control
- Sempra may damage or delay establishment of newly seeded, sodded or sprigged turf that is not well established.
- Sempra may damage or delay recovery of turf weakened by weather conditions or by physical damage such as scarification, coring, aeration or top dressing.

## Withholding Period

- Do not apply clippings from treated areas for mulch around fruit trees or vegetables
- Do not feed treated grass clippings to poultry or other livestock
- Do not allow grazing of treated turf.



## Sempra Compatibility

Sempra is compatible with the following herbicides commonly used in turf:

Kamba M or other dicamba/MCPA formulations;  
Bromicide MA or other bromoxynil/MCPA formulations: Amicide 625, Amicide 625 Low Odour or other amine formulations of 2,4-D; Roundup, Roundup CT

## For more information contact your Nufarm Regional Office:

Northern Region QLD/NSW (07) 3893 8777.

Southern Region VIC (03) 9282 1238, SA (08) 8444 6300.

Western Region WA (08) 9411 4000.

This publication is a guide only and no substitute for professional or expert advice. The product label should be consulted before use of any of the products referred to in this publication. Nufarm Australia Limited shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.

© Copyright 2004.

Nufarm Australia Limited

A.C.N. 004 377 780